

Middle-Four Maps and Net Categories

BRIAN DAY *

April 2, 2009

Abstract

We briefly relate the existence of a middle-four interchange map in a category with two monoidal structures, to the standard Cockett and Seely notion of a weakly distributive category.

1 Introduction

From a middle-four map we shall obtain an example of a structure (on the category of $I_{\oplus}$ -bimodules) with two weak distributivity maps:

$$\begin{aligned} d^l : A \otimes (B \oplus C) &\longrightarrow (A \otimes B) \oplus C \\ d^r : (B \oplus C) \otimes A &\longrightarrow B \oplus (C \otimes A) \end{aligned}$$

satisfying the eleven (planar) conditions (2)-(4) and (7)-(14) of [1], but not necessarily [1] (1), (5), and (6). We call such a structure a “net” category (see [2]) with the view that it is a proof-net category ([2]) without negation, or, equivalently, a (planar) weakly distributive category [1] without the unit objects.

Moreover, we can talk about a middle-four category with negation, which satisfies four more suitable axioms (c.f. [1] §4 (15)-(18)), so that, on passing to the category of $I_{\oplus}$ -bimodules, we get a proof-net category with a middle-four map.

2 Middle-Four Categories

First, we consider a category V with two monoidal structures $(\otimes, I_{\otimes}, a, l, r)$ and $(\oplus, I_{\oplus})$, where we have assumed that the second structure is strictly monoidal. Suppose this category is equipped with a natural middle-four transformation:

$$m : (A \oplus B) \otimes (C \oplus D) \longrightarrow (A \otimes C) \oplus (B \otimes D)$$

*Department of Mathematics, Macquarie University, NSW 2109, Australia.

and maps:

$$\begin{aligned}\mu : I_{\oplus} \otimes I_{\oplus} &\longrightarrow I_{\oplus} \\ \eta : I_{\otimes} &\longrightarrow I_{\oplus}\end{aligned}$$

satisfying the following four axioms:

M1

$$(a \oplus a)m(m \otimes 1) = m(1 \otimes m)a : ((U \oplus V) \otimes (W \oplus X)) \otimes (Y \oplus Z) \longrightarrow (U \otimes (W \otimes Y)) \oplus (V \otimes (X \otimes Z))$$

M2

$$(m \oplus 1)m = (1 \oplus m)m$$

from

$$(U \oplus V \oplus W) \otimes (X \oplus Y \oplus Z)$$

to

$$(U \otimes X) \oplus (V \otimes Y) \oplus (W \oplus Z)$$

and, if we put $I = I_{\otimes}$ and $R = I_{\oplus}$, we want (R, μ, η) to be a $\otimes$ -monoid in V such that both

M3

$$(1 \oplus (\mu \otimes 1))m(m \otimes 1) = m$$

from

$$((A \oplus R) \otimes (B \oplus R)) \otimes (C \oplus D)$$

to

$$((A \otimes B) \otimes C) \oplus (R \otimes D)$$

M4

$$((1 \otimes \mu) \oplus 1)m(1 \otimes m) = m$$

from

$$(B \oplus C) \otimes ((R \oplus A) \otimes (R \oplus B))$$

to

$$(B \otimes R) \oplus (C \otimes (A \otimes D))$$

Moreover, a middle-four category with negation has maps γ and τ , where

$$\begin{aligned}\gamma : A \otimes A^{\perp} &\longrightarrow R \\ \tau : I &\longrightarrow A^{\perp} \oplus A\end{aligned}$$

satisfying four axioms similar to:

$$\begin{array}{ccc}
 (A \oplus R) \otimes I & \xrightarrow{1 \otimes \tau} & (A \oplus R) \otimes (A^\perp \oplus A) \\
 \downarrow r & & \downarrow m \\
 A & & (A \otimes A^\perp) \oplus (R \otimes A) \\
 \downarrow \eta \otimes 1 & & \downarrow \gamma \oplus 1 \\
 R \otimes A & \xleftarrow{=} & R \oplus (R \otimes A)
 \end{array}$$

commutes (cf. [1] §4).

3 The Category of R -bimodules as a net category

The category $V(R)$ of R -bimodules in V has two induced (coherent) tensor products. These are:

- (1) $A \otimes B$ with action:

$$\begin{aligned}
 R \otimes (A \otimes B) \otimes R &\xrightarrow{\sim} (R \otimes A) \otimes (B \otimes R) \\
 &\xrightarrow{\sim} (R \otimes A \otimes I) \otimes (I \otimes B \otimes R) \\
 &\xrightarrow{(1 \otimes \eta) \otimes (\eta \otimes 1)} (R \otimes A \otimes R) \otimes (R \otimes B \otimes R) \\
 &\xrightarrow{\alpha \otimes \beta} A \otimes B
 \end{aligned}$$

which is unitless, and

- (2) $A \oplus B$ with action

$$R \otimes (A \oplus B) \otimes R \xrightarrow{"m"} (R \otimes A \otimes R) \oplus (R \otimes B \otimes R) \xrightarrow{\alpha \oplus \beta} A \oplus B$$

which has the unit R .

Furthermore, it is easily checked that our basic transformation

$$m : (A \oplus B) \otimes (C \oplus D) \longrightarrow (A \otimes C) \oplus (B \otimes D)$$

becomes an R -bimodule homomorphism. Thus, on putting $B = R$ and $C = R$ respectively, and using the bimodule actions, we get two natural transformations:


$$\begin{aligned}
 d^l : A \otimes (B \oplus C) &\longrightarrow (A \otimes B) \oplus C \\
 d^r : (B \oplus C) \otimes A &\longrightarrow B \oplus (C \otimes A)
 \end{aligned}$$

of R -bimodules.

We now check some [1] axioms on d^l and d^r .

Axioms [1] (7) and (8) are trivial.

Axiom [1] (9):



where $\alpha(1 \otimes \alpha)a = \alpha(\mu \otimes 1)$ since D is an R -bimodule.

Axiom [1] (10) is similar, but uses M1 and M4.

Axiom [1] (11) uses M2 and the action $(*)$ of R on $A \oplus B$.

$$\begin{array}{ccc}
(A \oplus B \oplus C) \otimes D & \xrightarrow{d^r} & A \oplus ((B \oplus C) \otimes D) \\
\downarrow m & & \downarrow \alpha \oplus 1 \\
& \cdot & \\
& \downarrow 1 \oplus m & \\
& \cdot & \\
& \downarrow m \oplus 1 \quad \alpha \oplus 1 & \\
& \cdot & \\
& \downarrow \alpha \oplus \alpha \oplus 1 & \\
& \cdot & \\
& \downarrow \alpha \oplus 1 \quad 1 \oplus \alpha \oplus 1 & \\
A \oplus B \oplus (C \otimes D) & &
\end{array}$$

$M2$ (left side), $1 \oplus d^r$ (right side), $(*)$ (bottom center)

Axiom [1] (12) is similar and still uses M2, while Axiom [1](13) uses M2, and [1] (14) uses M1.

Thus, from any category with two monoidal structures related by a middle-four interchange transformation satisfying the standard axioms for such a map, we obtain an example of a net-category (that is, a proof-net category without negation) which is also equipped with a middle-four interchange map.

Communication on this article can be made through Micah McCurdy (Macquarie University, micah.mccurdy@gmail.com) who kindly typed this manuscript. I would also like to thank Ross Street for a helpful comment on the middle-four map.

References

- [1] Cockett, J.R.B. and R.A.G. Seely, *Weakly Distributive Categories*, Journal of Pure and Applied Algebra v114 n2 (1997).
- [2] Dosen, K., and Z. Petric, “Proof-net categories”, (Preprint) Math. Inst. Serbian Academy of Sciences and Arts (2005), arXiv:math/0503301v3